

Embeddings of vertex operator algebras associated to orthogonal affine Lie algebras

Ozren Perše

Abstract

Let $L_{D_\ell}(-\ell + \frac{3}{2}, 0)$ (resp. $L_{B_\ell}(-\ell + \frac{3}{2}, 0)$) be the simple vertex operator algebra associated to affine Lie algebra of type $D_\ell^{(1)}$ (resp. $B_\ell^{(1)}$) with the lowest admissible half-integer level $-\ell + \frac{3}{2}$. We show that $L_{D_\ell}(-\ell + \frac{3}{2}, 0)$ is a vertex subalgebra of $L_{B_\ell}(-\ell + \frac{3}{2}, 0)$ with the same conformal vector. For $\ell = 4$, $L_{D_4}(-\frac{5}{2}, 0)$ is a vertex subalgebra of three copies of $L_{B_4}(-\frac{5}{2}, 0)$ contained in $L_{F_4}(-\frac{5}{2}, 0)$, and all five of these vertex operator algebras have the same conformal vector.

1 Introduction

Let $\mathfrak{g}_{X_l}$ be the simple Lie algebra of type X_l , $\hat{\mathfrak{g}}_{X_l}$ the associated affine Lie algebra of type $X_l^{(1)}$, and $L_{X_l}(k, 0)$ the simple vertex operator algebra associated to $\hat{\mathfrak{g}}_{X_l}$ of level $k \in \mathbb{C}$, $k \neq -h^\vee$, with conformal vector obtained from the Segal-Sugawara construction. It is an interesting problem to find pairs of vertex operator algebras $L_{X_l}(k, 0)$ and $L_{Y_{l'}}(k', 0)$, such that $L_{Y_{l'}}(k', 0)$ is a vertex subalgebra of $L_{X_l}(k, 0)$ with the same conformal vector. In the case of positive integer levels, such examples are called conformal embeddings in mathematical physics (cf. [AGO], [BB], [SW]).

In [P2], we showed that $L_{B_4}(-\frac{5}{2}, 0)$ is a vertex subalgebra of $L_{F_4}(-\frac{5}{2}, 0)$ with the same conformal vector. The motivation for studying level $k = -\frac{5}{2}$ was that the corresponding central charge of these vertex operator algebras is the same (and equal to -20). In this case, one can also consider Lie

¹2000 *Mathematics Subject Classification*. Primary 17B69; Secondary 17B67, 81R10.

subalgebra $\mathfrak{g}_{D_4}$ of $\mathfrak{g}_{B_4}$ and the associated vertex operator algebra $L_{D_4}(-\frac{5}{2}, 0)$. It turns out that this vertex operator algebra also has central charge -20 .

More generally, one can consider the embedding of orthogonal Lie algebras $\mathfrak{g}_{D_l}$ into $\mathfrak{g}_{B_l}$, for any $l \geq 4$. On the level of affine Lie algebras and corresponding vertex operator algebras, this embedding has been studied for level $k = 1$ (cf. [FF]). Using explicit realizations of modules $L_{D_l}(1, 0)$ and $L_{B_l}(1, 0)$, one can see that $L_{D_l}(1, 0)$ is a vertex subalgebra of $L_{B_l}(1, 0)$, but the corresponding conformal vectors are different in this case.

In this paper we study a level k such that $L_{D_l}(k, 0)$ is a vertex subalgebra of $L_{B_l}(k, 0)$ with the same conformal vector. The equality of conformal vectors implies the equality of corresponding central charges

$$\frac{k \dim \mathfrak{g}_{D_l}}{k + h_{D_l}^\vee} = \frac{k \dim \mathfrak{g}_{B_l}}{k + h_{B_l}^\vee}$$

Apart from the trivial solution $k = 0$, the only solution of this equation is $k = -l + \frac{3}{2}$. This suggests the study of vertex operator algebras $L_{D_l}(-l + \frac{3}{2}, 0)$ and $L_{B_l}(-l + \frac{3}{2}, 0)$. Since there are no known realizations of these irreducible modules, we consider these modules as quotients of the corresponding generalized Verma modules.

Vertex operator algebra $L_{B_l}(-l + \frac{3}{2}, 0)$ has been studied in [P1]. The level $k = -l + \frac{3}{2}$ is an admissible level for $\hat{\mathfrak{g}}_{B_l}$, in the sense of Kac and Wakimoto (cf. [KW1], [KW2], [W]). The results on admissible modules from [KW1] and [KW2] imply that $L_{B_l}(-l + \frac{3}{2}, 0)$ is a quotient of the corresponding generalized Verma module modulo the ideal generated by one singular vector v_{B_l} of conformal weight 2. The explicit formula for that vector was determined in [P1], and is relatively simple (cf. relation (4.1)). Vertex operator algebras associated to affine Lie algebras with admissible levels have also recently been studied in [A1, A2, AM, DLM].

In this paper we study vertex operator algebra $L_{D_l}(-l + \frac{3}{2}, 0)$. We verify that level $k = -l + \frac{3}{2}$ is admissible for $\hat{\mathfrak{g}}_{D_l}$, and show (using [KW1], [KW2]) that $L_{D_l}(-l + \frac{3}{2}, 0)$ is a quotient of corresponding generalized Verma module modulo the ideal generated by one singular vector v_{D_l} of conformal weight 4. The formula for that singular vector is much more complicated than the formula for v_{B_l} (cf. relation (4.2)). Using explicit formulas for vectors v_{B_l} and v_{D_l} we show that the vector v_{D_l} is contained in the $\hat{\mathfrak{g}}_{B_l}$ -submodule generated by v_{B_l} of the corresponding generalized Verma module, which directly implies that $L_{D_l}(-l + \frac{3}{2}, 0)$ is a vertex subalgebra of $L_{B_l}(-l + \frac{3}{2}, 0)$. Using certain relations generated by vector v_{B_l} on the quotient $L_{B_l}(-l + \frac{3}{2}, 0)$ of generalized

Verma module, we show that the conformal vectors of $L_{D_l}(-l + \frac{3}{2}, 0)$ and $L_{B_l}(-l + \frac{3}{2}, 0)$ are the same.

In the special case $l = 4$, $L_{F_4}(-\frac{5}{2}, 0)$ contains three copies of $L_{B_4}(-\frac{5}{2}, 0)$ as vertex subalgebras with the same conformal vector (cf. [P2]). It follows that vertex operator algebra $L_{D_4}(-\frac{5}{2}, 0)$ is a vertex subalgebra of all three copies, with the same conformal vector.

2 Preliminaries

Let $(V, Y, \mathbf{1}, \omega)$ be a vertex operator algebra (cf. [Bo, FHL, FLM, LL]). A *vertex subalgebra* of vertex algebra V is a subspace U of V such that $\mathbf{1} \in U$ and $Y(a, z)U \subseteq U[[z, z^{-1}]]$ for any $a \in U$. Suppose that $(V, Y, \mathbf{1}, \omega)$ is a vertex operator algebra and $(U, Y, \mathbf{1}, \omega')$ a vertex subalgebra of V , that has a structure of vertex operator algebra. We say that U is a *vertex operator subalgebra* of V if $\omega' = \omega$.

Let $\mathfrak{g}$ be a simple Lie algebra over $\mathbb{C}$ with a triangular decomposition $\mathfrak{g} = \mathfrak{n}_- \oplus \mathfrak{h} \oplus \mathfrak{n}_+$. Let $(\cdot, \cdot) : \mathfrak{g} \times \mathfrak{g} \rightarrow \mathbb{C}$ be the Killing form, normalized by the condition $(\theta, \theta) = 2$, where θ is the highest root of $\mathfrak{g}$. The affine Lie algebra $\hat{\mathfrak{g}}$ associated to $\mathfrak{g}$ is the vector space $\mathfrak{g} \otimes \mathbb{C}[t, t^{-1}] \oplus \mathbb{C}c$ equipped with the usual bracket operation and the canonical central element c (cf. [K1]). Let $h^\vee$ be the dual Coxeter number of $\hat{\mathfrak{g}}$. Let $\hat{\mathfrak{g}} = \hat{\mathfrak{n}}_- \oplus \hat{\mathfrak{h}} \oplus \hat{\mathfrak{n}}_+$ be the corresponding triangular decomposition of $\hat{\mathfrak{g}}$.

For every weight $\lambda \in \hat{\mathfrak{h}}^*$, denote by $M(\lambda)$ the Verma module for $\hat{\mathfrak{g}}$ with highest weight λ , and by $L(\lambda)$ the irreducible $\hat{\mathfrak{g}}$ -module with highest weight λ . Let U be a $\mathfrak{g}$ -module, and let $k \in \mathbb{C}$. Denote by $N(k, U)$ the generalized Verma $\hat{\mathfrak{g}}$ -module of level k induced from U . For $\mu \in \mathfrak{h}^*$, denote by $V(\mu)$ the irreducible highest weight $\mathfrak{g}$ -module with highest weight μ . We shall use the notation $N(k, \mu)$ to denote the $\hat{\mathfrak{g}}$ -module $N(k, V(\mu))$. Denote by $J(k, \mu)$ the maximal proper submodule of $N(k, \mu)$ and by $L(k, \mu) = N(k, \mu)/J(k, \mu)$ the corresponding irreducible $\hat{\mathfrak{g}}$ -module.

Let $\hat{\Delta}^{\vee \text{re}}$ (resp. $\hat{\Delta}_+^{\vee \text{re}}$) $\subset \hat{\mathfrak{h}}$ be the set of real (resp. positive real) coroots of $\hat{\mathfrak{g}}$. Fix $\lambda \in \hat{\mathfrak{h}}^*$. Let $\hat{\Delta}_\lambda^{\vee \text{re}} = \{\alpha \in \hat{\Delta}^{\vee \text{re}} \mid \langle \lambda, \alpha \rangle \in \mathbb{Z}\}$, $\hat{\Delta}_{\lambda+}^{\vee \text{re}} = \hat{\Delta}_\lambda^{\vee \text{re}} \cap \hat{\Delta}_+^{\vee \text{re}}$, $\hat{\Pi}^\vee$ the set of simple coroots in $\hat{\Delta}^{\vee \text{re}}$ and $\hat{\Pi}_\lambda^\vee = \{\alpha \in \hat{\Delta}_{\lambda+}^{\vee \text{re}} \mid \alpha \text{ not equal to a sum of several coroots from } \hat{\Delta}_{\lambda+}^{\vee \text{re}}\}$. Recall that a weight $\lambda \in \hat{\mathfrak{h}}^*$ is called *admissible* (cf. [KW1], [KW2] and [W]) if the following properties are satisfied:

$$\langle \lambda + \rho, \alpha \rangle \notin -\mathbb{Z}_+ \text{ for all } \alpha \in \hat{\Delta}_+^{\vee \text{re}},$$

$$\mathbb{Q}\hat{\Delta}_\lambda^{\vee\text{re}} = \mathbb{Q}\hat{\Pi}^\vee.$$

We shall use the following result of V. Kac and M. Wakimoto:

Proposition 2.1 ([KW1, Corollary 2.1]) *Let λ be an admissible weight. Then*

$$L(\lambda) \cong \frac{M(\lambda)}{\sum_{\alpha \in \hat{\Pi}_\lambda^\vee} U(\hat{\mathfrak{g}})v^\alpha},$$

where $v^\alpha \in M(\lambda)$ is a singular vector of weight $r_\alpha \cdot \lambda$, the highest weight vector of $M(r_\alpha \cdot \lambda) = U(\hat{\mathfrak{g}})v^\alpha \subset M(\lambda)$.

For $k \neq -h^\vee$, $N(k, 0)$ has the structure of vertex operator algebra (cf. [FZ, FB, K2, LL, L]) with conformal vector

$$\omega = \frac{1}{2(k + h^\vee)} \sum_{i=1}^{\dim \mathfrak{g}} a^i(-1)b^i(-1)\mathbf{1}, \quad (2.1)$$

where $\{a^i\}_{i=1, \dots, \dim \mathfrak{g}}$ is an arbitrary basis of $\mathfrak{g}$, and $\{b^i\}_{i=1, \dots, \dim \mathfrak{g}}$ the corresponding dual basis of $\mathfrak{g}$ with respect to the form $(\cdot, \cdot)$.

Since every $\hat{\mathfrak{g}}$ -submodule of $N(k, 0)$ is also an ideal in the vertex operator algebra $N(k, 0)$, it follows that $L(k, 0)$ is a vertex operator algebra, for any $k \neq -h^\vee$.

3 Simple Lie algebras of type D_l and B_l

Let $l \geq 4$ and consider two l -dimensional vector spaces $A_1 = \oplus_{i=1}^l \mathbb{C}a_i$, $A_2 = \oplus_{i=1}^l \mathbb{C}a_i^*$. Let $A = A_1 \oplus A_2$. The Clifford algebra $\text{Cliff}(A)$ is the associative algebra over $\mathbb{C}$ generated by A and relations

$$[a_i, a_j]_+ = [a_i^*, a_j^*]_+ = 0, \quad [a_i, a_j^*]_+ = \delta_{ij}, \quad i, j = 1, \dots, l.$$

The normal ordering on A is defined by

$$:xy: = \frac{1}{2}(xy - yx), \quad x, y \in A.$$

Then (cf. [Bou] and [FF]) all normally ordered quadratic elements $:xy:$ span a Lie algebra $\mathfrak{g}_{D_l}$ of type D_l with Cartan subalgebra $\mathfrak{h}$ spanned by

$$H_i = :a_i a_i^*: , \quad i = 1, \dots, l.$$

Let $\{\epsilon_i \mid 1 \leq i \leq l\} \subset \mathfrak{h}^*$ be the dual basis such that $\epsilon_i(H_j) = \delta_{ij}$. The root system of $\mathfrak{g}_{D_l}$ is given by

$$\Delta_{D_l} = \{\pm(\epsilon_i \pm \epsilon_j) \mid 1 \leq i, j \leq l, i < j\}$$

with $\alpha_1 = \epsilon_1 - \epsilon_2, \dots, \alpha_{l-1} = \epsilon_{l-1} - \epsilon_l, \alpha_l = \epsilon_{l-1} + \epsilon_l$ being a set of simple roots. The highest root is $\theta = \epsilon_1 + \epsilon_2$. The set of positive roots is $\Delta_{D_l}^+ = \{\epsilon_i \pm \epsilon_j \mid i < j\}$. We fix the root vectors

$$e_{\epsilon_i - \epsilon_j} = :a_i a_j^*:, \quad e_{\epsilon_i + \epsilon_j} = :a_i a_j:, \quad f_{\epsilon_i - \epsilon_j} = :a_j a_i^*:, \quad f_{\epsilon_i + \epsilon_j} = :a_j^* a_i^*:,$$

for $i, j = 1, \dots, l$ and $i < j$.

If the linear space A is added on to quadratic elements $:xy:, x, y \in A$, one obtains the simple Lie algebra $\mathfrak{g}_{B_l}$ of type B_l . The root system of $\mathfrak{g}_{B_l}$ with respect to the same Cartan subalgebra $\mathfrak{h}$ is

$$\Delta_{B_l} = \Delta_{D_l} \cup \{\pm\epsilon_i \mid 1 \leq i \leq l\}$$

with $\beta_1 = \epsilon_1 - \epsilon_2, \dots, \beta_{l-1} = \epsilon_{l-1} - \epsilon_l, \beta_l = \epsilon_l$ being a set of simple roots. The set of positive roots is $\Delta_{B_l}^+ = \Delta_{D_l}^+ \cup \{\epsilon_i \mid 1 \leq i \leq l\}$. Fix the root vectors for short roots

$$e_{\epsilon_i} = a_i, \quad f_{\epsilon_i} = a_i^*, \quad i = 1, \dots, l.$$

Denote by $h_\alpha = \alpha^\vee = [e_\alpha, f_\alpha]$ coroots, for any positive root $\alpha \in \Delta_{B_l}^+$. Clearly $h_{\epsilon_i} = 2H_i$, for $i = 1, \dots, l$.

4 Vertex operator algebras $L_{B_l}(-l + \frac{3}{2}, 0)$ and $L_{D_l}(-l + \frac{3}{2}, 0)$

Let $\hat{\mathfrak{g}}_{D_l}$ and $\hat{\mathfrak{g}}_{B_l}$ be affine Lie algebras associated to $\mathfrak{g}_{D_l}$ and $\mathfrak{g}_{B_l}$, respectively. For $k \in \mathbb{C}$, denote by $N_{D_l}(k, 0)$ and $N_{B_l}(k, 0)$ generalized Verma modules associated to $\hat{\mathfrak{g}}_{D_l}$ and $\hat{\mathfrak{g}}_{B_l}$ of level k , and by $L_{D_l}(k, 0)$ and $L_{B_l}(k, 0)$ corresponding irreducible modules.

We shall also use the notation $M_{D_l}(\lambda)$ (resp. $M_{B_l}(\lambda)$) for the Verma module for $\hat{\mathfrak{g}}_{D_l}$ (resp. $\hat{\mathfrak{g}}_{B_l}$) with highest weight $\lambda \in \hat{\mathfrak{h}}^*$, and $L_{D_l}(\lambda)$ (resp. $L_{B_l}(\lambda)$) for the irreducible $\hat{\mathfrak{g}}_{D_l}$ -module (resp. $\hat{\mathfrak{g}}_{B_l}$ -module) with highest weight $\lambda \in \hat{\mathfrak{h}}^*$.

Since $\mathfrak{g}_{D_l}$ is a Lie subalgebra of $\mathfrak{g}_{B_l}$, it follows that $\hat{\mathfrak{g}}_{D_l}$ is a Lie subalgebra of $\hat{\mathfrak{g}}_{B_l}$. Using the P-B-W theorem (which gives an embedding of the universal

enveloping algebra of a Lie subalgebra into the universal enveloping algebra of a Lie algebra) we obtain an embedding of generalized Verma module $N_{D_l}(k, 0)$ into $N_{B_l}(k, 0)$. Therefore, $N_{D_l}(k, 0)$ is a vertex subalgebra of $N_{B_l}(k, 0)$.

4.1 Vertex operator algebra $L_{B_l}(-l + \frac{3}{2}, 0)$

In this subsection we recall some facts about vertex operator algebra $L_{B_l}(-l + \frac{3}{2}, 0)$. It was proved in [P1, Theorem 11] that the maximal $\hat{\mathfrak{g}}_{B_l}$ -submodule of $N_{B_l}(-l + \frac{3}{2}, 0)$ is generated by one singular vector:

Proposition 4.1 *The maximal $\hat{\mathfrak{g}}_{B_l}$ -submodule of $N_{B_l}(-l + \frac{3}{2}, 0)$ is $J_{B_l}(-l + \frac{3}{2}, 0) = U(\hat{\mathfrak{g}}_{B_l})v_{B_l}$, where*

$$v_{B_l} = -\frac{1}{4}e_{\epsilon_1}(-1)^2\mathbf{1} + \sum_{j=2}^l e_{\epsilon_1 - \epsilon_j}(-1)e_{\epsilon_1 + \epsilon_j}(-1)\mathbf{1} \quad (4.1)$$

is a singular vector in $N_{B_l}(-l + \frac{3}{2}, 0)$.

Thus

$$L_{B_l}(-l + \frac{3}{2}, 0) \cong \frac{N_{B_l}(-l + \frac{3}{2}, 0)}{U(\hat{\mathfrak{g}}_{B_l})v_{B_l}}.$$

4.2 Vertex operator algebra $L_{D_l}(-l + \frac{3}{2}, 0)$

In this subsection we show that the maximal $\hat{\mathfrak{g}}_{D_l}$ -submodule of $N_{D_l}(-l + \frac{3}{2}, 0)$ is generated by one singular vector. We need two lemmas to prove that.

Denote by λ the weight $(-l + \frac{3}{2})\Lambda_0$. Then $N_{D_l}(-l + \frac{3}{2}, 0)$ is a quotient of the Verma module $M_{D_l}(\lambda)$ and $L_{D_l}(-l + \frac{3}{2}, 0) \cong L_{D_l}(\lambda)$.

Lemma 4.2 *The weight $\lambda = (-l + \frac{3}{2})\Lambda_0$ is admissible for $\hat{\mathfrak{g}}_{D_l}$ and*

$$\hat{\Pi}_\lambda^\vee = \{(2\delta - \theta)^\vee, \alpha_1^\vee, \dots, \alpha_l^\vee\}.$$

Proof: Clearly

$$\begin{aligned} \langle \lambda + \rho, \alpha_i^\vee \rangle &= 1 \quad \text{for } 1 \leq i \leq l, \\ \langle \lambda + \rho, \alpha_0^\vee \rangle &= -l + \frac{5}{2}, \end{aligned}$$

which implies

$$\langle \lambda + \rho, (2\delta - \theta)^\vee \rangle = \langle \lambda + \rho, 2\alpha_0^\vee + \alpha_1^\vee + 2\alpha_2^\vee + \dots 2\alpha_{l-2}^\vee + \alpha_{l-1}^\vee + \alpha_l^\vee \rangle = 2.$$

The claim of lemma now follows easily. $\square$

Lemma 4.3 *Vector*

$$\begin{aligned} v_{D_l} &= \frac{2(2l+1)}{3} \sum_{i=3}^l \sum_{\substack{j=3 \\ j \neq i}}^l e_{\epsilon_1 - \epsilon_i}(-1) e_{\epsilon_1 + \epsilon_i}(-1) e_{\epsilon_2 - \epsilon_j}(-1) e_{\epsilon_2 + \epsilon_j}(-1) \mathbf{1} \\ &+ \frac{2l+1}{3} \sum_{i=3}^l e_{\epsilon_1 - \epsilon_i}(-1) e_{\epsilon_1 + \epsilon_i}(-1) e_{\epsilon_2 - \epsilon_i}(-1) e_{\epsilon_2 + \epsilon_i}(-1) \mathbf{1} \\ &- \frac{2l+1}{3} \sum_{i=3}^l \sum_{\substack{j=3 \\ j \neq i}}^l e_{\epsilon_1 - \epsilon_i}(-1) e_{\epsilon_2 + \epsilon_i}(-1) e_{\epsilon_1 + \epsilon_j}(-1) e_{\epsilon_2 - \epsilon_j}(-1) \mathbf{1} \\ &- \frac{2l+1}{6} \sum_{i=3}^l \sum_{j=3}^l e_{\epsilon_1 - \epsilon_i}(-1) e_{\epsilon_2 + \epsilon_i}(-1) e_{\epsilon_1 - \epsilon_j}(-1) e_{\epsilon_2 + \epsilon_j}(-1) \mathbf{1} \\ &- \frac{2l+1}{6} \sum_{i=3}^l \sum_{j=3}^l e_{\epsilon_1 + \epsilon_i}(-1) e_{\epsilon_2 - \epsilon_i}(-1) e_{\epsilon_1 + \epsilon_j}(-1) e_{\epsilon_2 - \epsilon_j}(-1) \mathbf{1} \\ &+ 2 \sum_{i=3}^l \sum_{j=3}^{i-1} e_{\epsilon_1 + \epsilon_2}(-1) e_{\epsilon_1 + \epsilon_j}(-1) e_{\epsilon_2 - \epsilon_i}(-1) f_{\epsilon_j - \epsilon_i}(-1) \mathbf{1} \\ &+ 2 \sum_{i=3}^l \sum_{j=i+1}^l e_{\epsilon_1 + \epsilon_2}(-1) e_{\epsilon_1 + \epsilon_j}(-1) e_{\epsilon_2 - \epsilon_i}(-1) e_{\epsilon_i - \epsilon_j}(-1) \mathbf{1} \\ &- 2 \sum_{i=3}^l \sum_{j=3}^{i-1} e_{\epsilon_1 + \epsilon_2}(-1) e_{\epsilon_1 - \epsilon_j}(-1) e_{\epsilon_2 - \epsilon_i}(-1) e_{\epsilon_j + \epsilon_i}(-1) \mathbf{1} \\ &+ 2 \sum_{i=3}^l \sum_{j=i+1}^l e_{\epsilon_1 + \epsilon_2}(-1) e_{\epsilon_1 - \epsilon_j}(-1) e_{\epsilon_2 - \epsilon_i}(-1) e_{\epsilon_i + \epsilon_j}(-1) \mathbf{1} \\ &+ 2 \sum_{i=3}^l \sum_{j=3}^{i-1} e_{\epsilon_1 + \epsilon_2}(-1) e_{\epsilon_2 + \epsilon_i}(-1) e_{\epsilon_1 + \epsilon_j}(-1) f_{\epsilon_j + \epsilon_i}(-1) \mathbf{1} \\ &- 2 \sum_{i=3}^l \sum_{j=i+1}^l e_{\epsilon_1 + \epsilon_2}(-1) e_{\epsilon_2 + \epsilon_i}(-1) e_{\epsilon_1 + \epsilon_j}(-1) f_{\epsilon_i + \epsilon_j}(-1) \mathbf{1} \end{aligned}$$

$$\begin{aligned}
& -2 \sum_{i=3}^l \sum_{j=3}^{i-1} e_{\epsilon_1+\epsilon_2}(-1) e_{\epsilon_2+\epsilon_i}(-1) e_{\epsilon_1-\epsilon_j}(-1) e_{\epsilon_j-\epsilon_i}(-1) \mathbf{1} \\
& -2 \sum_{i=3}^l \sum_{j=i+1}^l e_{\epsilon_1+\epsilon_2}(-1) e_{\epsilon_2+\epsilon_i}(-1) e_{\epsilon_1-\epsilon_j}(-1) f_{\epsilon_i-\epsilon_j}(-1) \mathbf{1} \\
& -\frac{2(2l-5)}{3} \sum_{i=3}^l e_{\epsilon_1+\epsilon_2}(-1) f_{\epsilon_1-\epsilon_2}(-1) e_{\epsilon_1-\epsilon_i}(-1) e_{\epsilon_1+\epsilon_i}(-1) \mathbf{1} \\
& +\frac{2(2l-5)}{3} \sum_{i=3}^l e_{\epsilon_1+\epsilon_2}(-1) e_{\epsilon_1-\epsilon_2}(-1) e_{\epsilon_2-\epsilon_i}(-1) e_{\epsilon_2+\epsilon_i}(-1) \mathbf{1} \\
& +\sum_{i=3}^l e_{\epsilon_1+\epsilon_2}(-1) e_{\epsilon_1+\epsilon_i}(-1) e_{\epsilon_2-\epsilon_i}(-1) h_{\epsilon_i}(-1) \mathbf{1} \\
& +\frac{2l-5}{3} \sum_{i=3}^l e_{\epsilon_1+\epsilon_2}(-1) e_{\epsilon_1+\epsilon_i}(-1) e_{\epsilon_2-\epsilon_i}(-1) h_{\epsilon_1-\epsilon_2}(-1) \mathbf{1} \\
& -\sum_{i=3}^l e_{\epsilon_1+\epsilon_2}(-1) e_{\epsilon_1-\epsilon_i}(-1) e_{\epsilon_2+\epsilon_i}(-1) h_{\epsilon_i}(-1) \mathbf{1} \\
& +\frac{2l-5}{3} \sum_{i=3}^l e_{\epsilon_1+\epsilon_2}(-1) e_{\epsilon_1-\epsilon_i}(-1) e_{\epsilon_2+\epsilon_i}(-1) h_{\epsilon_1-\epsilon_2}(-1) \mathbf{1} \\
& +\frac{2l-5}{3} \sum_{i=3}^l e_{\epsilon_1+\epsilon_2}(-1) e_{\epsilon_1+\epsilon_i}(-2) e_{\epsilon_2-\epsilon_i}(-1) \mathbf{1} \\
& +\frac{2l-5}{3} \sum_{i=3}^l e_{\epsilon_1+\epsilon_2}(-1) e_{\epsilon_1-\epsilon_i}(-2) e_{\epsilon_2+\epsilon_i}(-1) \mathbf{1} \\
& -\frac{(2l+1)(l-3)}{3} \sum_{i=3}^l e_{\epsilon_1+\epsilon_2}(-2) e_{\epsilon_1+\epsilon_i}(-1) e_{\epsilon_2-\epsilon_i}(-1) \mathbf{1} \\
& -\frac{(2l+1)(l-3)}{3} \sum_{i=3}^l e_{\epsilon_1+\epsilon_2}(-2) e_{\epsilon_1-\epsilon_i}(-1) e_{\epsilon_2+\epsilon_i}(-1) \mathbf{1} \\
& -\frac{2l-5}{3} \sum_{i=3}^l e_{\epsilon_1+\epsilon_2}(-1) e_{\epsilon_1+\epsilon_i}(-1) e_{\epsilon_2-\epsilon_i}(-2) \mathbf{1}
\end{aligned}$$

$$\begin{aligned}
& -\frac{2l-5}{3} \sum_{i=3}^l e_{\epsilon_1+\epsilon_2}(-1)e_{\epsilon_1-\epsilon_i}(-1)e_{\epsilon_2+\epsilon_i}(-2)\mathbf{1} \\
& + \frac{(2l-5)(2l-1)}{6} e_{\epsilon_1+\epsilon_2}(-2)e_{\epsilon_1+\epsilon_2}(-1)h_{\epsilon_1-\epsilon_2}(-1)\mathbf{1} \\
& - \frac{2l-5}{2} e_{\epsilon_1+\epsilon_2}(-2)e_{\epsilon_1+\epsilon_2}(-1)h_{\epsilon_1}(-1)\mathbf{1} \\
& - \frac{(2l+1)(2l-5)(2l-7)}{24} e_{\epsilon_1+\epsilon_2}(-2)^2\mathbf{1} + \frac{(2l-5)^2}{2} e_{\epsilon_1+\epsilon_2}(-1)e_{\epsilon_1+\epsilon_2}(-3)\mathbf{1} \\
& - \frac{2(2l-5)(l-2)}{3(2l-1)} e_{\epsilon_1+\epsilon_2}(-1)^2(e_{\epsilon_1-\epsilon_2}(-1)f_{\epsilon_1-\epsilon_2}(-1) + f_{\epsilon_1-\epsilon_2}(-1)e_{\epsilon_1-\epsilon_2}(-1))\mathbf{1} \\
& + \frac{2l-5}{2l-1} \sum_{i=3}^l e_{\epsilon_1+\epsilon_2}(-1)^2(e_{\epsilon_2-\epsilon_i}(-1)f_{\epsilon_2-\epsilon_i}(-1) + f_{\epsilon_2-\epsilon_i}(-1)e_{\epsilon_2-\epsilon_i}(-1))\mathbf{1} \\
& + \frac{2l-5}{2l-1} \sum_{i=3}^l e_{\epsilon_1+\epsilon_2}(-1)^2(e_{\epsilon_2+\epsilon_i}(-1)f_{\epsilon_2+\epsilon_i}(-1) + f_{\epsilon_2+\epsilon_i}(-1)e_{\epsilon_2+\epsilon_i}(-1))\mathbf{1} \\
& + \frac{2l-5}{2l-1} \sum_{i=3}^l e_{\epsilon_1+\epsilon_2}(-1)^2(e_{\epsilon_1-\epsilon_i}(-1)f_{\epsilon_1-\epsilon_i}(-1) + f_{\epsilon_1-\epsilon_i}(-1)e_{\epsilon_1-\epsilon_i}(-1))\mathbf{1} \\
& + \frac{2l-5}{2l-1} \sum_{i=3}^l e_{\epsilon_1+\epsilon_2}(-1)^2(e_{\epsilon_1+\epsilon_i}(-1)f_{\epsilon_1+\epsilon_i}(-1) + f_{\epsilon_1+\epsilon_i}(-1)e_{\epsilon_1+\epsilon_i}(-1))\mathbf{1} \\
& + \frac{2l-5}{2l-1} e_{\epsilon_1+\epsilon_2}(-1)^2(e_{\epsilon_1+\epsilon_2}(-1)f_{\epsilon_1+\epsilon_2}(-1) + f_{\epsilon_1+\epsilon_2}(-1)e_{\epsilon_1+\epsilon_2}(-1))\mathbf{1} \\
& - \frac{4}{2l-1} \sum_{\substack{\alpha \in \Delta_{D_l}^+ \\ (\alpha, \epsilon_1)=0, (\alpha, \epsilon_2)=0}} e_{\epsilon_1+\epsilon_2}(-1)^2(e_{\alpha}(-1)f_{\alpha}(-1)\mathbf{1} + f_{\alpha}(-1)e_{\alpha}(-1))\mathbf{1} \\
& + e_{\epsilon_1+\epsilon_2}(-1)^2 \left(-\frac{2l-5}{6} h_{\epsilon_1-\epsilon_2}(-1)^2\mathbf{1} - \frac{1}{2(2l-1)} h_{\epsilon_1}(-1)^2\mathbf{1} \right. \\
& \left. - \frac{2l-5}{2l-1} h_{\epsilon_1-\epsilon_2}(-1)h_{\epsilon_1+\epsilon_2}(-1)\mathbf{1} + \frac{4}{2l-1} \sum_{i=3}^l h_{\epsilon_1-\epsilon_i}(-1)h_{\epsilon_1+\epsilon_i}(-1)\mathbf{1} \right) \\
& + \frac{2l-5}{2} e_{\epsilon_1+\epsilon_2}(-1)^2 h_{\epsilon_1+\epsilon_2}(-2)\mathbf{1} \tag{4.2}
\end{aligned}$$

is a singular vector in $N_{D_l}(-l + \frac{3}{2}, 0)$.

Proof: Direct verification of relations $e_{\epsilon_k-\epsilon_{k+1}}(0).v_{D_l} = 0$, for $k = 1, \dots, l-1$,

$e_{\epsilon_{l-1}+\epsilon_l}(0).v_{D_l} = 0$ and $f_\theta(1).v_{D_l} = 0$ (see Appendix). $\square$

Theorem 4.4 *The maximal $\hat{\mathfrak{g}}_{D_l}$ -submodule of $N_{D_l}(-l + \frac{3}{2}, 0)$ is $J_{D_l}(-l + \frac{3}{2}, 0) = U(\hat{\mathfrak{g}}_{D_l})v_{D_l}$, where v_{D_l} is given by relation (4.2).*

Proof: It follows from Proposition 2.1 and Lemma 4.2 that the maximal submodule of the Verma module $M_{D_l}(\lambda)$ is generated by $l+1$ singular vectors with weights

$$r_{2\delta-\theta}.\lambda, r_{\alpha_1}.\lambda, \dots, r_{\alpha_l}.\lambda.$$

It follows from Lemma 4.3 that v_{D_l} is a singular vector with weight $\lambda - 4\delta + 2\theta = r_{2\delta-\theta}.\lambda$. Other singular vectors have weights

$$r_{\alpha_i}.\lambda = \lambda - \langle \lambda + \rho, \alpha_i^\vee \rangle \alpha_i = \lambda - \alpha_i, \quad 1 \leq i \leq l,$$

so the images of these vectors under the projection of $M_{D_l}(\lambda)$ onto $N_{D_l}(-l + \frac{3}{2}, 0)$ are 0. Therefore, the maximal submodule of $N_{D_l}(-l + \frac{3}{2}, 0)$ is generated by the vector v_{D_l} , i.e. $J_{D_l}(-l + \frac{3}{2}, 0) = U(\hat{\mathfrak{g}}_{D_l})v_{D_l}$. $\square$

It follows that

$$L_{D_l}(-l + \frac{3}{2}, 0) \cong \frac{N_{D_l}(-l + \frac{3}{2}, 0)}{U(\hat{\mathfrak{g}}_{D_l})v_{D_l}}.$$

5 Embedding of $L_{D_l}(-l + \frac{3}{2}, 0)$ into $L_{B_l}(-l + \frac{3}{2}, 0)$

In this section we show that vector v_{D_l} is in the $\hat{\mathfrak{g}}_{B_l}$ -submodule $J_{B_l}(-l + \frac{3}{2}, 0)$ of $N_{B_l}(-l + \frac{3}{2}, 0)$ generated by vector v_{B_l} :

Lemma 5.1 $v_{D_l} \in J_{B_l}(-l + \frac{3}{2}, 0)$

Proof: Using relation (4.1), one can show that the following relations hold in $J_{B_l}(-l + \frac{3}{2}, 0)$:

$$\begin{aligned} f_{\epsilon_1}(0).v_{B_l} &= \frac{1}{2}e_{\epsilon_1}(-1)h_{\epsilon_1}(-1)\mathbf{1} + \sum_{j=2}^l e_{\epsilon_1+\epsilon_j}(-1)f_{\epsilon_j}(-1)\mathbf{1} + \sum_{j=2}^l e_{\epsilon_1-\epsilon_j}(-1)e_{\epsilon_j}(-1)\mathbf{1} \\ &\quad - \frac{2l-3}{2}e_{\epsilon_1}(-2)\mathbf{1}, \\ f_{\epsilon_1-\epsilon_i}(0).v_{B_l} &= -\frac{1}{2}e_{\epsilon_1}(-1)e_{\epsilon_i}(-1)\mathbf{1} + \sum_{j=2}^{i-1} e_{\epsilon_1+\epsilon_j}(-1)f_{\epsilon_j-\epsilon_i}(-1)\mathbf{1} \end{aligned} \tag{5.1}$$

$$\begin{aligned}
& - \sum_{j=2}^{i-1} e_{\epsilon_1-\epsilon_j}(-1)e_{\epsilon_j+\epsilon_i}(-1)\mathbf{1} - h_{\epsilon_1-\epsilon_i}(-1)e_{\epsilon_1+\epsilon_i}(-1)\mathbf{1} + \sum_{j=i+1}^l e_{\epsilon_1+\epsilon_j}(-1)e_{\epsilon_i-\epsilon_j}(-1)\mathbf{1} \\
& + \sum_{j=i+1}^l e_{\epsilon_1-\epsilon_j}(-1)e_{\epsilon_i+\epsilon_j}(-1)\mathbf{1} + \frac{2l-3}{2}e_{\epsilon_1+\epsilon_i}(-2)\mathbf{1}, \quad \text{for } i = 2, \dots, l,
\end{aligned} \tag{5.2}$$

$$\begin{aligned}
f_{\epsilon_1}(0)f_{\epsilon_1-\epsilon_2}(0).v_{B_l} &= \frac{1}{2}e_{\epsilon_2}(-1)h_{\epsilon_2}(-1)\mathbf{1} - e_{\epsilon_1+\epsilon_2}(-1)f_{\epsilon_1}(-1)\mathbf{1} + e_{\epsilon_1}(-1)f_{\epsilon_1-\epsilon_2}(-1)\mathbf{1} \\
& + \sum_{j=3}^l e_{\epsilon_2-\epsilon_j}(-1)e_{\epsilon_j}(-1)\mathbf{1} + \sum_{j=3}^l e_{\epsilon_2+\epsilon_j}(-1)f_{\epsilon_j}(-1)\mathbf{1} - \frac{2l-5}{2}e_{\epsilon_2}(-2)\mathbf{1},
\end{aligned} \tag{5.3}$$

$$\begin{aligned}
f_{\epsilon_1}(0)^2.v_{B_l} &= \frac{1}{2}(e_{\epsilon_1}(-1)f_{\epsilon_1}(-1)\mathbf{1} + f_{\epsilon_1}(-1)e_{\epsilon_1}(-1)\mathbf{1}) - \frac{1}{2}h_{\epsilon_1}(-1)^2\mathbf{1} \\
& + \sum_{j=2}^l (e_{\epsilon_j}(-1)f_{\epsilon_j}(-1)\mathbf{1} + f_{\epsilon_j}(-1)e_{\epsilon_j}(-1)\mathbf{1}) \\
& - \sum_{j=2}^l (e_{\epsilon_1+\epsilon_j}(-1)f_{\epsilon_1+\epsilon_j}(-1)\mathbf{1} + f_{\epsilon_1+\epsilon_j}(-1)e_{\epsilon_1+\epsilon_j}(-1)\mathbf{1}) \\
& - \sum_{j=2}^l (e_{\epsilon_1-\epsilon_j}(-1)f_{\epsilon_1-\epsilon_j}(-1)\mathbf{1} + e_{\epsilon_1-\epsilon_j}(-1)f_{\epsilon_1-\epsilon_j}(-1)\mathbf{1})
\end{aligned} \tag{5.4}$$

$$\begin{aligned}
f_{\epsilon_1-\epsilon_i}(0)f_{\epsilon_1+\epsilon_i}(0).v_{B_l} &= -\frac{1}{4}(e_{\epsilon_i}(-1)f_{\epsilon_i}(-1)\mathbf{1} + f_{\epsilon_i}(-1)e_{\epsilon_i}(-1)\mathbf{1}) \\
& + \frac{1}{4}(e_{\epsilon_1}(-1)f_{\epsilon_1}(-1)\mathbf{1} + f_{\epsilon_1}(-1)e_{\epsilon_1}(-1)\mathbf{1}) + h_{\epsilon_1-\epsilon_i}(-1)h_{\epsilon_1+\epsilon_i}(-1)\mathbf{1} \\
& + \frac{1}{2} \sum_{\substack{j=2 \\ j \neq i}}^l (e_{\epsilon_1+\epsilon_j}(-1)f_{\epsilon_1+\epsilon_j}(-1)\mathbf{1} + f_{\epsilon_1+\epsilon_j}(-1)e_{\epsilon_1+\epsilon_j}(-1)\mathbf{1}) \\
& + \frac{1}{2} \sum_{\substack{j=2 \\ j \neq i}}^l (e_{\epsilon_1-\epsilon_j}(-1)f_{\epsilon_1-\epsilon_j}(-1)\mathbf{1} + f_{\epsilon_1-\epsilon_j}(-1)e_{\epsilon_1-\epsilon_j}(-1)\mathbf{1}) \\
& - \frac{1}{2} \sum_{j=2}^{i-1} (e_{\epsilon_j-\epsilon_i}(-1)f_{\epsilon_j-\epsilon_i}(-1)\mathbf{1} + f_{\epsilon_j-\epsilon_i}(-1)e_{\epsilon_j-\epsilon_i}(-1)\mathbf{1}) \\
& - \frac{1}{2} \sum_{\substack{j=2 \\ j \neq i}}^l (e_{\epsilon_i+\epsilon_j}(-1)f_{\epsilon_i+\epsilon_j}(-1)\mathbf{1} + f_{\epsilon_i+\epsilon_j}(-1)e_{\epsilon_i+\epsilon_j}(-1)\mathbf{1})
\end{aligned}$$

$$-\frac{1}{2} \sum_{j=i+1}^l (e_{\epsilon_i-\epsilon_j}(-1)f_{\epsilon_i-\epsilon_j}(-1)\mathbf{1} + f_{\epsilon_i-\epsilon_j}(-1)e_{\epsilon_i-\epsilon_j}(-1)\mathbf{1}), \quad \text{for } i = 2, \dots, l, \quad (5.5)$$

$$\begin{aligned} f_{\epsilon_1+\epsilon_i}(0).v_{B_l} &= -\frac{1}{2}e_{\epsilon_1}(-1)f_{\epsilon_i}(-1)\mathbf{1} + \sum_{j=2}^{i-1} e_{\epsilon_1+\epsilon_j}(-1)f_{\epsilon_j+\epsilon_i}(-1)\mathbf{1} \\ &\quad - \sum_{j=2}^{i-1} e_{\epsilon_1-\epsilon_j}(-1)e_{\epsilon_j-\epsilon_i}(-1)\mathbf{1} - h_{\epsilon_1+\epsilon_i}(-1)e_{\epsilon_1-\epsilon_i}(-1)\mathbf{1} - \sum_{j=i+1}^l e_{\epsilon_1+\epsilon_j}(-1)f_{\epsilon_i+\epsilon_j}(-1)\mathbf{1} \\ &\quad - \sum_{j=i+1}^l e_{\epsilon_1-\epsilon_j}(-1)f_{\epsilon_i-\epsilon_j}(-1)\mathbf{1} + \frac{2l-3}{2}e_{\epsilon_1-\epsilon_i}(-2)\mathbf{1}, \quad \text{for } i = 3, \dots, l, \end{aligned} \quad (5.6)$$

$$\begin{aligned} f_{\epsilon_1-\epsilon_i}(0)f_{\epsilon_1-\epsilon_2}(0).v_{B_l} &= -\frac{1}{2}e_{\epsilon_2}(-1)e_{\epsilon_i}(-1)\mathbf{1} - h_{\epsilon_2-\epsilon_i}(-1)e_{\epsilon_2+\epsilon_i}(-1)\mathbf{1} \\ &\quad - f_{\epsilon_1-\epsilon_2}(-1)e_{\epsilon_1+\epsilon_i}(-1)\mathbf{1} - e_{\epsilon_1+\epsilon_2}(-1)f_{\epsilon_1-\epsilon_i}(-1)\mathbf{1} + \sum_{j=3}^{i-1} e_{\epsilon_2+\epsilon_j}(-1)f_{\epsilon_j-\epsilon_i}(-1)\mathbf{1} \\ &\quad - \sum_{j=3}^{i-1} e_{\epsilon_2-\epsilon_j}(-1)e_{\epsilon_i+\epsilon_j}(-1)\mathbf{1} + \sum_{j=i+1}^l e_{\epsilon_2-\epsilon_j}(-1)e_{\epsilon_i+\epsilon_j}(-1)\mathbf{1} \\ &\quad + \sum_{j=i+1}^l e_{\epsilon_2+\epsilon_j}(-1)e_{\epsilon_i-\epsilon_j}(-1)\mathbf{1} + \frac{2l-3}{2}e_{\epsilon_2+\epsilon_i}(-2)\mathbf{1}, \quad \text{for } i = 3, \dots, l, \end{aligned} \quad (5.7)$$

$$f_{\epsilon_1-\epsilon_2}(0)^2.v_{B_l} = -\frac{1}{2}e_{\epsilon_2}(-1)^2\mathbf{1} - 2f_{\epsilon_1-\epsilon_2}(-1)e_{\epsilon_1+\epsilon_2}(-1)\mathbf{1} + 2 \sum_{j=3}^l e_{\epsilon_2-\epsilon_j}(-1)e_{\epsilon_2+\epsilon_j}(-1)\mathbf{1}, \quad (5.8)$$

$$\begin{aligned} f_{\epsilon_1-\epsilon_2}(0)f_{\epsilon_1+\epsilon_i}(0).v_{B_l} &= -\frac{1}{2}e_{\epsilon_2}(-1)f_{\epsilon_i}(-1)\mathbf{1} - h_{\epsilon_2+\epsilon_i}(-1)e_{\epsilon_2-\epsilon_i}(-1)\mathbf{1} \\ &\quad - f_{\epsilon_1-\epsilon_2}(-1)e_{\epsilon_1-\epsilon_i}(-1)\mathbf{1} - e_{\epsilon_1+\epsilon_2}(-1)f_{\epsilon_1+\epsilon_i}(-1)\mathbf{1} + \sum_{j=3}^{i-1} e_{\epsilon_2+\epsilon_j}(-1)f_{\epsilon_i+\epsilon_j}(-1)\mathbf{1} \\ &\quad - \sum_{j=3}^{i-1} e_{\epsilon_2-\epsilon_j}(-1)e_{\epsilon_j-\epsilon_i}(-1)\mathbf{1} - \sum_{j=i+1}^l e_{\epsilon_2-\epsilon_j}(-1)f_{\epsilon_i-\epsilon_j}(-1)\mathbf{1} \\ &\quad - \sum_{j=i+1}^l e_{\epsilon_2+\epsilon_j}(-1)f_{\epsilon_i+\epsilon_j}(-1)\mathbf{1} + \frac{2l-3}{2}e_{\epsilon_2-\epsilon_i}(-2)\mathbf{1}, \quad \text{for } i = 3, \dots, l. \end{aligned} \quad (5.9)$$

Using relations (5.1)–(5.9) one can obtain

$$\begin{aligned}
v_{D_l} = & \left(\frac{2l+1}{12} e_{\epsilon_2}(-1)^2 - \frac{2l-5}{3} f_{\epsilon_1-\epsilon_2}(-1) e_{\epsilon_1+\epsilon_2}(-1) \right. \\
& + \frac{2l+1}{3} \sum_{i=3}^l e_{\epsilon_2-\epsilon_i}(-1) e_{\epsilon_2+\epsilon_i}(-1) - \frac{1}{2} e_{\epsilon_1+\epsilon_2}(-1) e_{\epsilon_2}(-1) f_{\epsilon_1}(0) \\
& + \sum_{i=3}^l e_{\epsilon_1+\epsilon_2}(-1) e_{\epsilon_2-\epsilon_i}(-1) f_{\epsilon_1-\epsilon_i}(0) - \frac{2l+1}{12} e_{\epsilon_1}(-1) e_{\epsilon_2}(-1) f_{\epsilon_1-\epsilon_2}(0) \\
& - \frac{(2l+1)(2l-5)}{12} e_{\epsilon_1+\epsilon_2}(-2) f_{\epsilon_1-\epsilon_2}(0) + \frac{2l-5}{6} h_{\epsilon_1-\epsilon_2}(-1) e_{\epsilon_1+\epsilon_2}(-1) f_{\epsilon_1-\epsilon_2}(0) \\
& - \frac{2l+1}{6} \sum_{i=3}^l e_{\epsilon_1-\epsilon_i}(-1) e_{\epsilon_2+\epsilon_i}(-1) f_{\epsilon_1-\epsilon_2}(0) \\
& - \frac{2l+1}{6} \sum_{i=3}^l e_{\epsilon_1+\epsilon_i}(-1) e_{\epsilon_2-\epsilon_i}(-1) f_{\epsilon_1-\epsilon_2}(0) + \frac{1}{2} e_{\epsilon_1}(-1) e_{\epsilon_1+\epsilon_2}(-1) f_{\epsilon_1}(0) f_{\epsilon_1-\epsilon_2}(0) \\
& + \frac{1}{2l-1} e_{\epsilon_1+\epsilon_2}(-1)^2 f_{\epsilon_1}(0)^2 - \frac{2l-5}{2l-1} e_{\epsilon_1+\epsilon_2}(-1)^2 f_{\epsilon_1-\epsilon_2}(0) f_{\epsilon_1+\epsilon_2}(0) \\
& + \frac{4}{2l-1} \sum_{i=3}^l e_{\epsilon_1+\epsilon_2}(-1)^2 f_{\epsilon_1-\epsilon_i}(0) f_{\epsilon_1+\epsilon_i}(0) \\
& + \sum_{i=3}^l e_{\epsilon_1+\epsilon_2}(-1) e_{\epsilon_2+\epsilon_i}(-1) f_{\epsilon_1+\epsilon_i}(0) - \sum_{i=3}^l e_{\epsilon_1+\epsilon_2}(-1) e_{\epsilon_1-\epsilon_i}(-1) f_{\epsilon_1-\epsilon_i}(0) f_{\epsilon_1-\epsilon_2}(0) \\
& + \frac{2l+1}{24} e_{\epsilon_1}(-1)^2 f_{\epsilon_1-\epsilon_2}(0)^2 + \frac{2l-5}{6} e_{\epsilon_1-\epsilon_2}(-1) e_{\epsilon_1+\epsilon_2}(-1) f_{\epsilon_1-\epsilon_2}(0)^2 \\
& + \frac{2l+1}{6} \sum_{i=3}^l e_{\epsilon_1-\epsilon_i}(-1) e_{\epsilon_1+\epsilon_i}(-1) f_{\epsilon_1-\epsilon_2}(0)^2 \\
& \left. - \sum_{i=3}^l e_{\epsilon_1+\epsilon_2}(-1) e_{\epsilon_1+\epsilon_i}(-1) f_{\epsilon_1-\epsilon_2}(0) f_{\epsilon_1+\epsilon_i}(0) \right) \cdot v_{B_l},
\end{aligned}$$

which implies the claim of lemma. $\square$

It follows from Lemma 5.1, Theorem 4.4 and Proposition 4.1 that the embedding of $N_{D_l}(-l + \frac{3}{2}, 0)$ into $N_{B_l}(-l + \frac{3}{2}, 0)$ induces the embedding of $L_{D_l}(-l + \frac{3}{2}, 0)$ into $L_{B_l}(-l + \frac{3}{2}, 0)$. We get:

Theorem 5.2 $L_{D_l}(-l + \frac{3}{2}, 0)$ is a vertex subalgebra of $L_{B_l}(-l + \frac{3}{2}, 0)$.

Next, we show that $L_{D_l}(-l + \frac{3}{2}, 0)$ is a vertex operator subalgebra of $L_{B_l}(-l + \frac{3}{2}, 0)$, i.e. that these vertex operator algebras have the same conformal vector. We use the following lemma:

Lemma 5.3 *Relation*

$$\begin{aligned} & (2l-1) \sum_{\substack{\alpha \in \Delta_{B_l}^+ \\ (\alpha, \alpha)=1}} (e_\alpha(-1)f_\alpha(-1)\mathbf{1} + f_\alpha(-1)e_\alpha(-1)\mathbf{1}) \\ &= 4 \sum_{\alpha \in \Delta_{D_l}^+} (e_\alpha(-1)f_\alpha(-1)\mathbf{1} + f_\alpha(-1)e_\alpha(-1)\mathbf{1}) + \sum_{\substack{\alpha \in \Delta_{B_l}^+ \\ (\alpha, \alpha)=1}} h_\alpha(-1)^2 \mathbf{1} \end{aligned}$$

holds in $L_{B_l}(-l + \frac{3}{2}, 0)$.

Proof: Proposition 4.1 implies that relation

$$\left(2lf_{\epsilon_1}(0)^2 + 4 \sum_{i=2}^l f_{\epsilon_1 - \epsilon_i}(0)f_{\epsilon_1 + \epsilon_i}(0) \right) \cdot v_{B_l} = 0$$

holds in $L_{B_l}(-l + \frac{3}{2}, 0)$. Using relations (5.4) and (5.5) we get the claim of lemma. $\square$

Theorem 5.4 *Denote by ω_{B_l} the conformal vector for vertex operator algebra $L_{B_l}(-l + \frac{3}{2}, 0)$, and by ω_{D_l} the conformal vector for $L_{D_l}(-l + \frac{3}{2}, 0)$. Then*

$$\omega_{B_l} = \omega_{D_l}.$$

Proof: Set $\{\frac{1}{2}h_\alpha \mid \alpha \in \Delta_{B_l}^+, (\alpha, \alpha) = 1\}$ is an orthonormal basis of $\mathfrak{h}$ with respect to the form $(\cdot, \cdot)$. Lemma 5.3 implies that

$$\begin{aligned} & \sum_{\substack{\alpha \in \Delta_{B_l}^+ \\ (\alpha, \alpha)=1}} (e_\alpha(-1)f_\alpha(-1)\mathbf{1} + f_\alpha(-1)e_\alpha(-1)\mathbf{1}) \\ &= \frac{4}{2l-1} \sum_{\alpha \in \Delta_{D_l}^+} (e_\alpha(-1)f_\alpha(-1)\mathbf{1} + f_\alpha(-1)e_\alpha(-1)\mathbf{1}) + \frac{1}{2l-1} \sum_{\substack{\alpha \in \Delta_{B_l}^+ \\ (\alpha, \alpha)=1}} h_\alpha(-1)^2 \mathbf{1} \end{aligned} \tag{5.10}$$

holds in $L_{B_l}(-l + \frac{3}{2}, 0)$. It follows from relation (2.1) that

$$\begin{aligned}
\omega_{B_l} &= \frac{1}{2l+1} \left(\frac{1}{4} \sum_{\substack{\alpha \in \Delta_{B_l}^+ \\ (\alpha, \alpha)=1}} h_\alpha(-1)^2 \mathbf{1} + \sum_{\alpha \in \Delta_{B_l}^+} \frac{(\alpha, \alpha)}{2} (e_\alpha(-1)f_\alpha(-1)\mathbf{1} + f_\alpha(-1)e_\alpha(-1)\mathbf{1}) \right) \\
&= \frac{1}{2l+1} \left(\frac{1}{4} \sum_{\substack{\alpha \in \Delta_{B_l}^+ \\ (\alpha, \alpha)=1}} h_\alpha(-1)^2 \mathbf{1} + \sum_{\alpha \in \Delta_{D_l}^+} (e_\alpha(-1)f_\alpha(-1)\mathbf{1} + f_\alpha(-1)e_\alpha(-1)\mathbf{1}) \right. \\
&\quad \left. + \frac{1}{2} \sum_{\substack{\alpha \in \Delta_{B_l}^+ \\ (\alpha, \alpha)=1}} (e_\alpha(-1)f_\alpha(-1) + f_\alpha(-1)e_\alpha(-1)\mathbf{1}) \right).
\end{aligned}$$

Using relation (5.10), we obtain

$$\omega_{B_l} = \frac{1}{2l-1} \left(\frac{1}{4} \sum_{\substack{\alpha \in \Delta_{B_l}^+ \\ (\alpha, \alpha)=1}} h_\alpha(-1)^2 \mathbf{1} + \sum_{\alpha \in \Delta_{D_l}^+} (e_\alpha(-1)f_\alpha(-1)\mathbf{1} + f_\alpha(-1)e_\alpha(-1)\mathbf{1}) \right) = \omega_{D_l}. \quad \square$$

Thus, $L_{D_l}(-l + \frac{3}{2}, 0)$ is a vertex operator subalgebra of $L_{B_l}(-l + \frac{3}{2}, 0)$.

6 Case $l = 4$

For $l = 4$, simple Lie algebra $\mathfrak{g}_{F_4}$ contains three copies of $\mathfrak{g}_{B_4}$ as Lie subalgebras. We denote these copies $\mathfrak{g}_{B_4}$, $\mathfrak{g}'_{B_4}$ and $\mathfrak{g}''_{B_4}$, and we have the corresponding isomorphisms

$$\begin{aligned}
\pi' : \mathfrak{g}_{B_4} &\rightarrow \mathfrak{g}'_{B_4}, \\
\pi'' : \mathfrak{g}_{B_4} &\rightarrow \mathfrak{g}''_{B_4}.
\end{aligned}$$

Lie algebra $\mathfrak{g}_{D_4}$ is a Lie subalgebra of all three copies of $\mathfrak{g}_{B_4}$, and it is invariant under isomorphisms π' and π'' . When restricted to $\mathfrak{g}_{D_4}$, π' and π'' are automorphisms induced from the automorphisms of the root system Δ_{D_4} , determined by:

$$\begin{aligned}
\pi'(\alpha_1) &= \alpha_3, \quad \pi'(\alpha_2) = \alpha_2, \quad \pi'(\alpha_3) = \alpha_4, \quad \pi'(\alpha_4) = \alpha_1 \\
\pi''(\alpha_1) &= \alpha_4, \quad \pi''(\alpha_2) = \alpha_2, \quad \pi''(\alpha_3) = \alpha_3, \quad \pi''(\alpha_4) = \alpha_1.
\end{aligned} \tag{6.1}$$

Vertex operator algebra $L_{F_4}(-\frac{5}{2}, 0)$ contains three copies of $L_{B_4}(-\frac{5}{2}, 0)$ as vertex operator subalgebras (cf. [P2]). We denote these copies $L_{B_4}(-\frac{5}{2}, 0)$, $L'_{B_4}(-\frac{5}{2}, 0)$ and $L''_{B_4}(-\frac{5}{2}, 0)$, and the corresponding isomorphisms of vertex operator algebras are induced from π' and π'' . Using relations (6.1) and (4.2), one can easily verify that $\pi'(v_{D_4}) = v_{D_4}$ and $\pi''(v_{D_4}) = v_{D_4}$, which implies that $L_{D_4}(-\frac{5}{2}, 0)$ is a vertex subalgebra of all three copies $L_{B_4}(-\frac{5}{2}, 0)$, $L'_{B_4}(-\frac{5}{2}, 0)$ and $L''_{B_4}(-\frac{5}{2}, 0)$. We obtain:

Corollary 6.1 *$L_{D_4}(-\frac{5}{2}, 0)$ is a vertex subalgebra of three copies of $L_{B_4}(-\frac{5}{2}, 0)$ contained in $L_{F_4}(-\frac{5}{2}, 0)$. Vertex operator algebras $L_{D_4}(-\frac{5}{2}, 0)$, $L_{B_4}(-\frac{5}{2}, 0)$, $L'_{B_4}(-\frac{5}{2}, 0)$, $L''_{B_4}(-\frac{5}{2}, 0)$ and $L_{F_4}(-\frac{5}{2}, 0)$ have the same conformal vector.*

7 Appendix

In this Appendix we show relation $e_{\epsilon_1-\epsilon_2}(0).v_{D_l} = 0$ from the proof of Lemma 4.3. Relations $e_{\epsilon_2-\epsilon_3}(0).v_{D_l} = 0$, $e_{\epsilon_k-\epsilon_{k+1}}(0).v_{D_l} = 0$, for $k = 3, \dots, l-1$, $e_{\epsilon_{l-1}+\epsilon_l}(0).v_{D_l} = 0$ and $f_\theta(1).v_{D_l} = 0$ can be verified similarly. We have:

$$\begin{aligned} & e_{\epsilon_1-\epsilon_2}(0).e_{\epsilon_1-\epsilon_i}(-1)e_{\epsilon_1+\epsilon_i}(-1)e_{\epsilon_2-\epsilon_j}(-1)e_{\epsilon_2+\epsilon_j}(-1)\mathbf{1} \\ &= e_{\epsilon_1-\epsilon_i}(-1)e_{\epsilon_1+\epsilon_i}(-1)e_{\epsilon_1-\epsilon_j}(-1)e_{\epsilon_2+\epsilon_j}(-1)\mathbf{1} \\ &+ e_{\epsilon_1-\epsilon_i}(-1)e_{\epsilon_1+\epsilon_i}(-1)e_{\epsilon_1+\epsilon_j}(-1)e_{\epsilon_2-\epsilon_j}(-1)\mathbf{1} \\ &+ e_{\epsilon_1-\epsilon_i}(-1)e_{\epsilon_1+\epsilon_i}(-1)e_{\epsilon_1+\epsilon_2}(-2)\mathbf{1}, \quad \text{for } i, j = 3, \dots, l, \end{aligned} \tag{7.1}$$

$$\begin{aligned} & e_{\epsilon_1-\epsilon_2}(0).e_{\epsilon_1-\epsilon_i}(-1)e_{\epsilon_2+\epsilon_i}(-1)e_{\epsilon_1+\epsilon_j}(-1)e_{\epsilon_2-\epsilon_j}(-1)\mathbf{1} \\ &= e_{\epsilon_1-\epsilon_i}(-1)e_{\epsilon_1+\epsilon_i}(-1)e_{\epsilon_1+\epsilon_j}(-1)e_{\epsilon_2-\epsilon_j}(-1)\mathbf{1} \\ &+ e_{\epsilon_1-\epsilon_i}(-1)e_{\epsilon_2+\epsilon_i}(-1)e_{\epsilon_1+\epsilon_j}(-1)e_{\epsilon_1-\epsilon_j}(-1)\mathbf{1}, \quad \text{for } i, j = 3, \dots, l, i \neq j, \end{aligned} \tag{7.2}$$

$$\begin{aligned} & e_{\epsilon_1-\epsilon_2}(0).e_{\epsilon_1-\epsilon_i}(-1)e_{\epsilon_2+\epsilon_i}(-1)e_{\epsilon_1-\epsilon_j}(-1)e_{\epsilon_2+\epsilon_j}(-1)\mathbf{1} \\ &= e_{\epsilon_1-\epsilon_i}(-1)e_{\epsilon_1+\epsilon_i}(-1)e_{\epsilon_1-\epsilon_j}(-1)e_{\epsilon_2+\epsilon_j}(-1)\mathbf{1} \\ &+ e_{\epsilon_1-\epsilon_i}(-1)e_{\epsilon_2+\epsilon_i}(-1)e_{\epsilon_1-\epsilon_j}(-1)e_{\epsilon_1+\epsilon_j}(-1)\mathbf{1}, \quad \text{for } i, j = 3, \dots, l, i \neq j, \end{aligned} \tag{7.3}$$

$$\begin{aligned} & e_{\epsilon_1-\epsilon_2}(0).e_{\epsilon_1-\epsilon_i}(-1)e_{\epsilon_2+\epsilon_i}(-1)e_{\epsilon_1-\epsilon_i}(-1)e_{\epsilon_2+\epsilon_i}(-1)\mathbf{1} \\ &= 2e_{\epsilon_1-\epsilon_i}(-1)^2e_{\epsilon_1+\epsilon_i}(-1)e_{\epsilon_2+\epsilon_i}(-1)\mathbf{1} + e_{\epsilon_1+\epsilon_2}(-2)e_{\epsilon_1-\epsilon_i}(-1)e_{\epsilon_1+\epsilon_i}(-1)\mathbf{1} \\ & \quad \text{for } i = 3, \dots, l, \end{aligned} \tag{7.4}$$

$$e_{\epsilon_1-\epsilon_2}(0).e_{\epsilon_1+\epsilon_i}(-1)e_{\epsilon_2-\epsilon_i}(-1)e_{\epsilon_1+\epsilon_j}(-1)e_{\epsilon_2-\epsilon_j}(-1)\mathbf{1}$$

$$\begin{aligned}
&= e_{\epsilon_1+\epsilon_i}(-1)e_{\epsilon_1-\epsilon_i}(-1)e_{\epsilon_1+\epsilon_j}(-1)e_{\epsilon_2-\epsilon_j}(-1)\mathbf{1} \\
&+ e_{\epsilon_1+\epsilon_i}(-1)e_{\epsilon_2-\epsilon_i}(-1)e_{\epsilon_1+\epsilon_j}(-1)e_{\epsilon_1-\epsilon_j}(-1)\mathbf{1}, \quad \text{for } i, j = 3, \dots, l, i \neq j,
\end{aligned} \tag{7.5}$$

$$\begin{aligned}
&e_{\epsilon_1-\epsilon_2}(0).e_{\epsilon_1+\epsilon_i}(-1)e_{\epsilon_2-\epsilon_i}(-1)e_{\epsilon_1+\epsilon_i}(-1)e_{\epsilon_2-\epsilon_i}(-1)\mathbf{1} \\
&= 2e_{\epsilon_1+\epsilon_i}(-1)^2e_{\epsilon_1-\epsilon_i}(-1)e_{\epsilon_2-\epsilon_i}(-1)\mathbf{1} + e_{\epsilon_1+\epsilon_2}(-2)e_{\epsilon_1+\epsilon_i}(-1)e_{\epsilon_1-\epsilon_i}(-1)\mathbf{1} \\
&\quad \text{for } i = 3, \dots, l,
\end{aligned} \tag{7.6}$$

$$\begin{aligned}
&e_{\epsilon_1-\epsilon_2}(0).e_{\epsilon_1+\epsilon_2}(-1)e_{\epsilon_1+\epsilon_j}(-1)e_{\epsilon_2-\epsilon_i}(-1)f_{\epsilon_j-\epsilon_i}(-1)\mathbf{1} \\
&= e_{\epsilon_1+\epsilon_2}(-1)e_{\epsilon_1+\epsilon_j}(-1)e_{\epsilon_1-\epsilon_i}(-1)f_{\epsilon_j-\epsilon_i}(-1)\mathbf{1}, \quad \text{for } i, j = 3, \dots, l, j < i,
\end{aligned} \tag{7.7}$$

$$\begin{aligned}
&e_{\epsilon_1-\epsilon_2}(0).e_{\epsilon_1+\epsilon_2}(-1)e_{\epsilon_1+\epsilon_j}(-1)e_{\epsilon_2-\epsilon_i}(-1)e_{\epsilon_i-\epsilon_j}(-1)\mathbf{1} \\
&= e_{\epsilon_1+\epsilon_2}(-1)e_{\epsilon_1+\epsilon_j}(-1)e_{\epsilon_1-\epsilon_i}(-1)e_{\epsilon_i-\epsilon_j}(-1)\mathbf{1} \quad \text{for } i, j = 3, \dots, l, i < j,
\end{aligned} \tag{7.8}$$

$$\begin{aligned}
&e_{\epsilon_1-\epsilon_2}(0).e_{\epsilon_1+\epsilon_2}(-1)e_{\epsilon_1-\epsilon_j}(-1)e_{\epsilon_2-\epsilon_i}(-1)e_{\epsilon_i+\epsilon_j}(-1)\mathbf{1} \\
&= e_{\epsilon_1+\epsilon_2}(-1)e_{\epsilon_1-\epsilon_j}(-1)e_{\epsilon_1-\epsilon_i}(-1)e_{\epsilon_i+\epsilon_j}(-1)\mathbf{1}, \quad \text{for } i, j = 3, \dots, l, i \neq j,
\end{aligned} \tag{7.9}$$

$$\begin{aligned}
&e_{\epsilon_1-\epsilon_2}(0).e_{\epsilon_1+\epsilon_2}(-1)e_{\epsilon_2+\epsilon_i}(-1)e_{\epsilon_1+\epsilon_j}(-1)f_{\epsilon_i+\epsilon_j}(-1)\mathbf{1} \\
&= e_{\epsilon_1+\epsilon_2}(-1)e_{\epsilon_1+\epsilon_i}(-1)e_{\epsilon_1+\epsilon_j}(-1)f_{\epsilon_i+\epsilon_j}(-1)\mathbf{1}, \quad \text{for } i, j = 3, \dots, l, i \neq j,
\end{aligned} \tag{7.10}$$

$$\begin{aligned}
&e_{\epsilon_1-\epsilon_2}(0).e_{\epsilon_1+\epsilon_2}(-1)e_{\epsilon_2+\epsilon_i}(-1)e_{\epsilon_1-\epsilon_j}(-1)e_{\epsilon_j-\epsilon_i}(-1)\mathbf{1} \\
&= e_{\epsilon_1+\epsilon_2}(-1)e_{\epsilon_1+\epsilon_i}(-1)e_{\epsilon_1-\epsilon_j}(-1)e_{\epsilon_j-\epsilon_i}(-1)\mathbf{1}, \quad \text{for } i, j = 3, \dots, l, j < i,
\end{aligned} \tag{7.11}$$

$$\begin{aligned}
&e_{\epsilon_1-\epsilon_2}(0).e_{\epsilon_1+\epsilon_2}(-1)e_{\epsilon_2+\epsilon_i}(-1)e_{\epsilon_1-\epsilon_j}(-1)f_{\epsilon_i-\epsilon_j}(-1)\mathbf{1} \\
&= e_{\epsilon_1+\epsilon_2}(-1)e_{\epsilon_1+\epsilon_i}(-1)e_{\epsilon_1-\epsilon_j}(-1)f_{\epsilon_i-\epsilon_j}(-1)\mathbf{1}, \quad \text{for } i, j = 3, \dots, l, i < j,
\end{aligned} \tag{7.12}$$

$$\begin{aligned}
&e_{\epsilon_1-\epsilon_2}(0).e_{\epsilon_1+\epsilon_2}(-1)f_{\epsilon_1-\epsilon_2}(-1)e_{\epsilon_1-\epsilon_i}(-1)e_{\epsilon_1+\epsilon_i}(-1)\mathbf{1} \\
&= e_{\epsilon_1+\epsilon_2}(-1)e_{\epsilon_1+\epsilon_i}(-1)e_{\epsilon_1-\epsilon_i}(-2)\mathbf{1} + e_{\epsilon_1+\epsilon_2}(-1)e_{\epsilon_1-\epsilon_i}(-1)e_{\epsilon_1+\epsilon_i}(-2)\mathbf{1} \\
&+ e_{\epsilon_1+\epsilon_2}(-1)e_{\epsilon_1-\epsilon_i}(-1)e_{\epsilon_1+\epsilon_i}(-1)h_{\epsilon_1-\epsilon_2}(-1)\mathbf{1}, \quad \text{for } i = 3, \dots, l,
\end{aligned} \tag{7.13}$$

$$\begin{aligned}
&e_{\epsilon_1-\epsilon_2}(0).e_{\epsilon_1+\epsilon_2}(-1)e_{\epsilon_1-\epsilon_2}(-1)e_{\epsilon_2-\epsilon_i}(-1)e_{\epsilon_2+\epsilon_i}(-1)\mathbf{1} \\
&= e_{\epsilon_1+\epsilon_2}(-1)e_{\epsilon_1-\epsilon_2}(-1)e_{\epsilon_1-\epsilon_i}(-1)e_{\epsilon_2+\epsilon_i}(-1)\mathbf{1} \\
&+ e_{\epsilon_1+\epsilon_2}(-1)e_{\epsilon_1-\epsilon_2}(-1)e_{\epsilon_1+\epsilon_2}(-2)\mathbf{1} \\
&+ e_{\epsilon_1+\epsilon_2}(-1)e_{\epsilon_1-\epsilon_2}(-1)e_{\epsilon_1+\epsilon_i}(-1)e_{\epsilon_2-\epsilon_i}(-1)\mathbf{1}, \quad \text{for } i = 3, \dots, l,
\end{aligned} \tag{7.14}$$

$$\begin{aligned} & e_{\epsilon_1 - \epsilon_2}(0) \cdot e_{\epsilon_1 + \epsilon_2}(-1) e_{\epsilon_1 + \epsilon_i}(-1) e_{\epsilon_2 - \epsilon_i}(-1) h_{\epsilon_i}(-1) \mathbf{1} \\ &= e_{\epsilon_1 + \epsilon_2}(-1) e_{\epsilon_1 + \epsilon_i}(-1) e_{\epsilon_1 - \epsilon_i}(-1) h_{\epsilon_i}(-1) \mathbf{1}, \quad \text{for } i = 3, \dots, l, \end{aligned} \quad (7.15)$$

$$\begin{aligned}
& e_{\epsilon_1-\epsilon_2}(0) \cdot e_{\epsilon_1+\epsilon_2}(-1) e_{\epsilon_1+\epsilon_i}(-1) e_{\epsilon_2-\epsilon_i}(-1) h_{\epsilon_1-\epsilon_2}(-1) \mathbf{1} \\
&= e_{\epsilon_1+\epsilon_2}(-1) e_{\epsilon_1+\epsilon_i}(-1) e_{\epsilon_1-\epsilon_i}(-1) h_{\epsilon_1-\epsilon_2}(-1) \mathbf{1} \\
&\quad - 2e_{\epsilon_1+\epsilon_2}(-1) e_{\epsilon_1-\epsilon_2}(-1) e_{\epsilon_1+\epsilon_i}(-1) e_{\epsilon_2-\epsilon_i}(-1) \mathbf{1} \\
&\quad + 2e_{\epsilon_1+\epsilon_2}(-1) e_{\epsilon_1+\epsilon_i}(-1) e_{\epsilon_1-\epsilon_i}(-2) \mathbf{1}, \quad \text{for } i = 3, \dots, l,
\end{aligned} \tag{7.16}$$

$$\begin{aligned} & e_{\epsilon_1 - \epsilon_2}(0) \cdot e_{\epsilon_1 + \epsilon_2}(-1) e_{\epsilon_1 - \epsilon_i}(-1) e_{\epsilon_2 + \epsilon_i}(-1) h_{\epsilon_i}(-1) \mathbf{1} \\ &= e_{\epsilon_1 + \epsilon_2}(-1) e_{\epsilon_1 - \epsilon_i}(-1) e_{\epsilon_1 + \epsilon_i}(-1) h_{\epsilon_i}(-1) \mathbf{1}, \quad \text{for } i = 3, \dots, l, \end{aligned} \quad (7.17)$$

$$\begin{aligned}
& e_{\epsilon_1-\epsilon_2}(0) \cdot e_{\epsilon_1+\epsilon_2}(-1) e_{\epsilon_1-\epsilon_i}(-1) e_{\epsilon_2+\epsilon_i}(-1) h_{\epsilon_1-\epsilon_2}(-1) \mathbf{1} \\
&= e_{\epsilon_1+\epsilon_2}(-1) e_{\epsilon_1-\epsilon_i}(-1) e_{\epsilon_1+\epsilon_i}(-1) h_{\epsilon_1-\epsilon_2}(-1) \mathbf{1} \\
&\quad - 2 e_{\epsilon_1+\epsilon_2}(-1) e_{\epsilon_1-\epsilon_2}(-1) e_{\epsilon_1-\epsilon_i}(-1) e_{\epsilon_2+\epsilon_i}(-1) \mathbf{1} \\
&\quad + 2 e_{\epsilon_1+\epsilon_2}(-1) e_{\epsilon_1-\epsilon_i}(-1) e_{\epsilon_1+\epsilon_i}(-2) \mathbf{1}, \quad \text{for } i = 3, \dots, l,
\end{aligned} \tag{7.18}$$

$$e_{\epsilon_1 - \epsilon_2}(0).e_{\epsilon_1 + \epsilon_2}(-1)e_{\epsilon_1 + \epsilon_i}(-2)e_{\epsilon_2 - \epsilon_i}(-1)\mathbf{1} = e_{\epsilon_1 + \epsilon_2}(-1)e_{\epsilon_1 + \epsilon_i}(-2)e_{\epsilon_1 - \epsilon_i}(-1)\mathbf{1}$$

$$\text{for } i = 3, \dots, l, \quad (7.19)$$

$$e_{\epsilon_1 - \epsilon_2}(0).e_{\epsilon_1 + \epsilon_2}(-1)e_{\epsilon_1 - \epsilon_i}(-2)e_{\epsilon_2 + \epsilon_i}(-1)\mathbf{1} = e_{\epsilon_1 + \epsilon_2}(-1)e_{\epsilon_1 - \epsilon_i}(-2)e_{\epsilon_1 + \epsilon_i}(-1)\mathbf{1}$$

for $i = 3, \dots, l$,

$$(7.20)$$

$$e_{\epsilon_1 - \epsilon_2}(0).e_{\epsilon_1 + \epsilon_2}(-2)e_{\epsilon_1 + \epsilon_i}(-1)e_{\epsilon_2 - \epsilon_i}(-1)\mathbf{1} = e_{\epsilon_1 + \epsilon_2}(-2)e_{\epsilon_1 + \epsilon_i}(-1)e_{\epsilon_1 - \epsilon_i}(-1)\mathbf{1}$$

$$\text{for } i = 3, \dots, l, \quad (7.21)$$

$$e_{\epsilon_1 - \epsilon_2}(0).e_{\epsilon_1 + \epsilon_2}(-2)e_{\epsilon_1 - \epsilon_i}(-1)e_{\epsilon_2 + \epsilon_i}(-1)\mathbf{1} = e_{\epsilon_1 + \epsilon_2}(-2)e_{\epsilon_1 - \epsilon_i}(-1)e_{\epsilon_1 + \epsilon_i}(-1)\mathbf{1}$$

for $i = 3, \dots, l$,

(7.22)

$$e_{\epsilon_1 - \epsilon_2}(0).e_{\epsilon_1 + \epsilon_2}(-1)e_{\epsilon_1 + \epsilon_i}(-1)e_{\epsilon_2 - \epsilon_i}(-2)\mathbf{1} = e_{\epsilon_1 + \epsilon_2}(-1)e_{\epsilon_1 + \epsilon_i}(-1)e_{\epsilon_1 - \epsilon_i}(-2)\mathbf{1}$$

for $i = 3, \dots, l$,

(7.23)

$$e_{\epsilon_1 - \epsilon_2}(0).e_{\epsilon_1 + \epsilon_2}(-1)e_{\epsilon_1 - \epsilon_i}(-1)e_{\epsilon_2 + \epsilon_i}(-2)\mathbf{1} = e_{\epsilon_1 + \epsilon_2}(-1)e_{\epsilon_1 - \epsilon_i}(-1)e_{\epsilon_1 + \epsilon_i}(-2)\mathbf{1}$$

for $i = 3, \dots, l$,

$$(7.24)$$

$$e_{\epsilon_1-\epsilon_2}(0).e_{\epsilon_1+\epsilon_2}(-2)e_{\epsilon_1+\epsilon_2}(-1)h_{\epsilon_1-\epsilon_2}(-1)\mathbf{1} = -2e_{\epsilon_1+\epsilon_2}(-2)e_{\epsilon_1+\epsilon_2}(-1)e_{\epsilon_1-\epsilon_2}(-1)\mathbf{1}, \quad (7.25)$$

$$e_{\epsilon_1 - \epsilon_2}(0).e_{\epsilon_1 + \epsilon_2}(-2)e_{\epsilon_1 + \epsilon_2}(-1)h_{\epsilon_1}(-1)\mathbf{1} = -2e_{\epsilon_1 + \epsilon_2}(-2)e_{\epsilon_1 + \epsilon_2}(-1)e_{\epsilon_1 - \epsilon_2}(-1)\mathbf{1}, \quad (7.26)$$

$$e_{\epsilon_1 - \epsilon_2}(0).e_{\epsilon_1 + \epsilon_2}(-2)^2 \mathbf{1} = 0, \quad (7.27)$$

$$e_{\epsilon_1 - \epsilon_2}(0).e_{\epsilon_1 + \epsilon_2}(-1)e_{\epsilon_1 + \epsilon_2}(-3)\mathbf{1} = 0, \quad (7.28)$$

$$e_{\epsilon_1-\epsilon_2}(0).e_{\epsilon_1+\epsilon_2}(-1)^2(e_{\epsilon_1-\epsilon_2}(-1)f_{\epsilon_1-\epsilon_2}(-1)+f_{\epsilon_1-\epsilon_2}(-1)e_{\epsilon_1-\epsilon_2}(-1))\mathbf{1}$$

$$= 2e_{\epsilon_1+\epsilon_2}(-1)^2 e_{\epsilon_1-\epsilon_2}(-1) h_{\epsilon_1-\epsilon_2}(-1) \mathbf{1} + 2e_{\epsilon_1+\epsilon_2}(-1)^2 e_{\epsilon_1-\epsilon_2}(-2) \mathbf{1}, \quad (7.29)$$

$$\begin{aligned} & e_{\epsilon_1-\epsilon_2}(0).e_{\epsilon_1+\epsilon_2}(-1)^2 (e_{\epsilon_2-\epsilon_i}(-1) f_{\epsilon_2-\epsilon_i}(-1) + f_{\epsilon_2-\epsilon_i}(-1) e_{\epsilon_2-\epsilon_i}(-1)) \mathbf{1} \\ &= 2e_{\epsilon_1+\epsilon_2}(-1)^2 e_{\epsilon_1-\epsilon_i}(-1) f_{\epsilon_2-\epsilon_i}(-1) \mathbf{1} - e_{\epsilon_1+\epsilon_2}(-1)^2 e_{\epsilon_1-\epsilon_2}(-2) \mathbf{1}, \\ & \text{for } i = 3, \dots, l, \end{aligned} \quad (7.30)$$

$$\begin{aligned} & e_{\epsilon_1-\epsilon_2}(0).e_{\epsilon_1+\epsilon_2}(-1)^2 (e_{\epsilon_2+\epsilon_i}(-1) f_{\epsilon_2+\epsilon_i}(-1) + f_{\epsilon_2+\epsilon_i}(-1) e_{\epsilon_2+\epsilon_i}(-1)) \mathbf{1} \\ &= 2e_{\epsilon_1+\epsilon_2}(-1)^2 e_{\epsilon_1+\epsilon_i}(-1) f_{\epsilon_2+\epsilon_i}(-1) \mathbf{1} - e_{\epsilon_1+\epsilon_2}(-1)^2 e_{\epsilon_1-\epsilon_2}(-2) \mathbf{1}, \\ & \text{for } i = 3, \dots, l, \end{aligned} \quad (7.31)$$

$$\begin{aligned} & e_{\epsilon_1-\epsilon_2}(0).e_{\epsilon_1+\epsilon_2}(-1)^2 (e_{\epsilon_1-\epsilon_i}(-1) f_{\epsilon_1-\epsilon_i}(-1) + f_{\epsilon_1-\epsilon_i}(-1) e_{\epsilon_1-\epsilon_i}(-1)) \mathbf{1} \\ &= -2e_{\epsilon_1+\epsilon_2}(-1)^2 e_{\epsilon_1-\epsilon_i}(-1) f_{\epsilon_2-\epsilon_i}(-1) \mathbf{1} + e_{\epsilon_1+\epsilon_2}(-1)^2 e_{\epsilon_1-\epsilon_2}(-2) \mathbf{1}, \\ & \text{for } i = 3, \dots, l, \end{aligned} \quad (7.32)$$

$$\begin{aligned} & e_{\epsilon_1-\epsilon_2}(0).e_{\epsilon_1+\epsilon_2}(-1)^2 (e_{\epsilon_1+\epsilon_i}(-1) f_{\epsilon_1+\epsilon_i}(-1) + f_{\epsilon_1+\epsilon_i}(-1) e_{\epsilon_1+\epsilon_i}(-1)) \mathbf{1} \\ &= -2e_{\epsilon_1+\epsilon_2}(-1)^2 e_{\epsilon_1+\epsilon_i}(-1) f_{\epsilon_2+\epsilon_i}(-1) \mathbf{1} + e_{\epsilon_1+\epsilon_2}(-1)^2 e_{\epsilon_1-\epsilon_2}(-2) \mathbf{1}, \\ & \text{for } i = 3, \dots, l, \end{aligned} \quad (7.33)$$

$$e_{\epsilon_1-\epsilon_2}(0).e_{\epsilon_1+\epsilon_2}(-1)^2 (e_{\epsilon_1+\epsilon_2}(-1) f_{\epsilon_1+\epsilon_2}(-1) + f_{\epsilon_1+\epsilon_2}(-1) e_{\epsilon_1+\epsilon_2}(-1)) \mathbf{1} = 0, \quad (7.34)$$

$$\begin{aligned} & e_{\epsilon_1-\epsilon_2}(0).e_{\epsilon_1+\epsilon_2}(-1)^2 (e_{\alpha}(-1) f_{\alpha}(-1) \mathbf{1} + f_{\alpha}(-1) e_{\alpha}(-1)) \mathbf{1} = 0, \\ & \text{for } \alpha \in \Delta_{D_l}^+ \text{ such that } (\alpha, \epsilon_1) = 0, (\alpha, \epsilon_2) = 0, \end{aligned} \quad (7.35)$$

$$\begin{aligned} & e_{\epsilon_1-\epsilon_2}(0).e_{\epsilon_1+\epsilon_2}(-1)^2 h_{\epsilon_1-\epsilon_2}(-1)^2 \mathbf{1} = -4e_{\epsilon_1+\epsilon_2}(-1)^2 e_{\epsilon_1-\epsilon_2}(-1) h_{\epsilon_1-\epsilon_2}(-1) \mathbf{1} \\ & -4e_{\epsilon_1+\epsilon_2}(-1)^2 e_{\epsilon_1-\epsilon_2}(-2) \mathbf{1}, \end{aligned} \quad (7.36)$$

$$\begin{aligned} & e_{\epsilon_1-\epsilon_2}(0).e_{\epsilon_1+\epsilon_2}(-1)^2 h_{\epsilon_1}(-1)^2 \mathbf{1} = -4e_{\epsilon_1+\epsilon_2}(-1)^2 e_{\epsilon_1-\epsilon_2}(-1) h_{\epsilon_1}(-1) \mathbf{1} \\ & -4e_{\epsilon_1+\epsilon_2}(-1)^2 e_{\epsilon_1-\epsilon_2}(-2) \mathbf{1}, \end{aligned} \quad (7.37)$$

$$\begin{aligned} & e_{\epsilon_1-\epsilon_2}(0).e_{\epsilon_1+\epsilon_2}(-1)^2 h_{\epsilon_1-\epsilon_2}(-1) h_{\epsilon_1+\epsilon_2}(-1) \mathbf{1} = -2e_{\epsilon_1+\epsilon_2}(-1)^2 e_{\epsilon_1-\epsilon_2}(-1) h_{\epsilon_1+\epsilon_2}(-1) \mathbf{1}, \\ & \end{aligned} \quad (7.38)$$

$$\begin{aligned} & e_{\epsilon_1-\epsilon_2}(0).e_{\epsilon_1+\epsilon_2}(-1)^2 h_{\epsilon_1-\epsilon_i}(-1) h_{\epsilon_1+\epsilon_i}(-1) \mathbf{1} = -e_{\epsilon_1+\epsilon_2}(-1)^2 e_{\epsilon_1-\epsilon_2}(-1) h_{\epsilon_1}(-1) \mathbf{1} \\ & -e_{\epsilon_1+\epsilon_2}(-1)^2 e_{\epsilon_1-\epsilon_2}(-2) \mathbf{1}, \quad \text{for } i = 3, \dots, l, \end{aligned} \quad (7.39)$$

$$e_{\epsilon_1-\epsilon_2}(0).e_{\epsilon_1+\epsilon_2}(-1)^2 h_{\epsilon_1+\epsilon_2}(-2) \mathbf{1} = 0. \quad (7.40)$$

Using relations (7.1)-(7.40) and formula (4.2) for vector v_{D_l} , one directly obtains $e_{\epsilon_1-\epsilon_2}(0).v_{D_l} = 0$.

References

- [A1] D. Adamović, Some rational vertex algebras, *Glas. Mat. Ser. III* **29(49)** (1994), 25–40.

- [A2] D. Adamović, A construction of admissible $A_1^{(1)}$ -modules of level $-\frac{4}{3}$, *J. Pure Appl. Algebra* **196** (2005), 119–134.
- [AM] D. Adamović and A. Milas, Vertex operator algebras associated to modular invariant representations for $A_1^{(1)}$, *Math. Res. Lett.* **2** (1995), 563–575.
- [AGO] R. C. Arcuri, J. F. Gomes and D. I. Olive, Conformal subalgebras and symmetric spaces, *Nuclear Phys. B* **285** (1987), 327–339.
- [BB] F. A. Bais and P. G. Bouwknegt, A classification of subgroup truncations of the bosonic string, *Nuclear Phys. B* **279** (1987), 561–570.
- [Bo] R. E. Borcherds, Vertex algebras, Kac-Moody algebras, and the Monster, *Proc. Natl. Acad. Sci. USA* **83** (1986), 3068–3071.
- [Bou] Bourbaki, Groupes et algèbres de Lie, Hermann, Paris, 1975
- [DLM] C. Dong, H. Li and G. Mason, Vertex operator algebras associated to admissible representations of $\widehat{\mathfrak{sl}}_2$, *Comm. Math. Phys.* **184** (1997), 65–93.
- [FF] A. J. Feingold and I. B. Frenkel, Classical affine algebras, *Adv. in Math.*, **56** (1985), 117–172.
- [FB] E. Frenkel and D. Ben-Zvi, *Vertex algebras and algebraic curves*, Mathematical Surveys and Monographs, 88, American Mathematical Society, Providence, RI, 2001.
- [FHL] I. Frenkel, Y.-Z. Huang and J. Lepowsky, On axiomatic approaches to vertex operator algebras and modules, *Mem. Amer. Math. Soc.* **104**, 1993.
- [FLM] I. Frenkel, J. Lepowsky and A. Meurman, *Vertex Operator Algebras and the Monster*, Pure and Appl. Math., **Vol. 134**, Academic Press, Boston, 1988.
- [FZ] I. Frenkel and Y.-C. Zhu, Vertex operator algebras associated to representations of affine and Virasoro algebras, *Duke Math. J.* **66** (1992), 123–168.

- [K1] V. G. Kac, *Infinite dimensional Lie algebras*, 3rd ed., Cambridge Univ. Press, Cambridge, 1990.
- [K2] V. G. Kac, *Vertex Algebras for Beginners*, University Lecture Series, Second Edition, AMS, Vol. 10, 1998.
- [KW1] V. Kac and M. Wakimoto, Modular invariant representations of infinite dimensional Lie algebras and superalgebras, *Proc. Natl. Acad. Sci. USA* **85** (1988), 4956–4960.
- [KW2] V. Kac and M. Wakimoto, Classification of modular invariant representations of affine algebras, in *Infinite Dimensional Lie algebras and groups*, Advanced Series in Math. Phys. **7**, World Scientific, Teaneck NJ, 1989.
- [LL] J. Lepowsky and H. Li, *Introduction to vertex operator algebras and their representations*, Progress in Math., **Vol. 227**, Birkhauser, Boston, 2004.
- [L] H.-S. Li, Local systems of vertex operators, vertex superalgebras and modules. *J. Pure Appl. Algebra* **109** (1996), 143–195.
- [P1] O. Perše, Vertex operator algebras associated to type B affine Lie algebras on admissible half-integer levels, *J. Algebra* **307** (2007), 215–248.
- [P2] O. Perše, Vertex operator algebra analogue of embedding of B_4 into F_4 , *J. Pure Appl. Algebra* **211** (2007), 702–720.
- [SW] A. N. Schellekens and N. P. Warner, Conformal subalgebras of Kac-Moody algebras, *Phys. Rev. D* **34** (1986), 3092–3096.
- [W] M. Wakimoto, *Lectures on infinite-dimensional Lie algebra*, World Scientific, River Edge NJ, 2001.

Department of Mathematics, University of Zagreb, Bijenička 30,
10000 Zagreb, Croatia
E-mail address: perse@math.hr